



Surface TB

HARD SURFACE DISINFECTANT



PHYSICAL PROPERTIES

Appearance: Clear Liquid

Odor: Citrus / floral

Ph (Conc.): 12-12.5

Flash Point: >212°F (>100°C) estimated.

PACKAGING

Quart, 2.5 gal, 4X1 gal, 5 gal

EPA REG. NO. 1839-83-74357 EPA EST. NO. 74357-MO-01

ABOUT THIS PRODUCT

This product is designed specifically as a general ready-to-use cleaner and disinfectant for use in homes, hospitals, nursing homes, patient rooms, operating rooms, ICU areas, shower rooms, locker rooms, public restrooms, schools, food processing plants, food service establishments, transportation terminals, office buildings, manufacturing facilities, lodging establishments, retail businesses, veterinary clinics, pet shops, animal life science laboratories and athlete/recreational facilities where housekeeping is of prime importance in controlling the hazard of cross contamination.

It is formulated to disinfect hard, non-porous, inanimate environmental surfaces: floors, walls, metal surfaces, stainless steel surfaces, glazed porcelain, glazed ceramic tile, plastic surfaces, chrome, brass, copper, laminated surfaces, baked enamel surfaces, bathrooms, shower stalls, bathtubs, and cabinets

For plastic and painted surfaces, spot test on an inconspicuous area before use. May be used in the kitchen on counters, sinks, appliances, and stovetops. A rinse with potable water is required for surfaces in direct contact with food.

DIRECTIONS

DISINFECTION, DEODORIZING AND CLEANING - Remove gross filth or heavy soil prior to application of the product. Hold container six to eight inches from surface to be treated. Spray area until it is covered with the solution. Allow product to penetrate and the surface to remain wet for 3 minutes (for Bloodborne Pathogens: allow surface to remain wet for HIV-1 1 minute and for HBV and HCV 5 minutes). No scrubbing is necessary. Wipe off with a clean cloth, mop or sponge. The product will not leave grit or soap scum.

To disinfect food processing premises: Before using this product food products and packaging materials must be removed or carefully protected. For floors, walls and storage areas, hold container six to eight inches from the surface to be treated and spray area until it is covered with the solution. Allow product to penetrate and remain wet for 3 minutes (for Bloodborne Pathogens: allow surface to remain wet for HIV-1 1 minute and for HBV and HCV 5 minutes). Wipe off with a clean cloth, mop or sponge and thoroughly rinse all surfaces in the area with potable water. Heavily soiled areas must be pre-cleaned before using this product.

FIRST AID

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person. If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice. NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.